

POTENTIALLY PHOTSENSITIZING MEDICATIONS

This is not an exhaustive list of all potentially photosensitizing drugs. The client is responsible for determining whether a medication photosensitizes themselves to the spectrum of light emitted from the PicoWay laser system.

Acne Meds

Isotretinoin (Accutane)
Tretinoin (Retin-A)

Anticancer

Chlorambucil
Cyclophosphamide
Dacarbazine
Fluorouracil
Flutamide
Mercaptopurine
Methotrexate
Procarbazine
Thioguanine
Vinblastine

Antidepressants

Amitriptyline
Amoxapine
Clomipramine
Doxepin
Imipramine
Isocarboxazid
Maprotiline
Phenelzine
Protriptyline
Trazadone
Trimipramine

Antiepileptics, Sedative, Muscle Relaxants

Carbamazepine
Cyclobenzaprine
Diazepam
Meprobamate
Phenobarbital
Phenytoin

Antihistamines

Azatadine
Clemastine
Diphenhydramine
Terfenadine
Tripelemnamine

Antihypertensives

Captopril
Diltiazem
Methyldopa
Minoxidil
Nifedipine

Antimicrobials

Ciprofloxacin
Clofazimine
Dapsone
Demeclocycline
Doxycycline
Enoxacin
Flucytosine
Griseofulvin
Ketoconazole
Lomefloxacin
Methacycline
Minocycline
Nalidixic acid
Narfloxacin
Ofloxacin
Oxytetracycline
Pyrazinamide
Sulfa drugs (Bactrim, Septra,
Tetracycline)

Antiparasitics

Bithionol
Chloroquine
Pyruvium
Pamoate
Quinine
Thiabendazole

Antipsychotics

Chlorpromazine
Chlorprothixene
Fluphenazine
Haloperidol
Perphenazine
Prochlorperazine
Promethazine
Thioridazine
Thiothixane
Trifluoperazine
Thioflupromazine
Trimeprazine

Cardiovascular

Amiodarone
Atenolol
Captopril
Diltiazem
Disopyramide
Nifedipine
Propranolol
Quinidine gluconate
Quinidine sulfate
Verapamil

Diuretics

Acetazolamide
Amiloride
Bendroflumethiazide
Benzthiazide
Chlorothiazide
Furosemide
Hydrochlorothiazide

Diuretics (cont.)

Hydro flumethiazide
Methyclothiazide
Metalazone
Polythiazide
Quinethazone
Trichlormethia-zide

Hypoglycemics

Acetohexamide
Chlorpropamide
Glipizide
Tolazamide Tolbutamide

NSAIDS

Diclofenac
Fenoprofen
Flurbiprofen
Indomethacin
Ketoprofen
Meclofenamate
Naproxen
Phenylbutazone
Piroxicam
Sulindac

Others

Bergamot oil
Oils of citron, lavender,
lime, sandalwood
Benzocaine
Clofibrate
Oral contraceptive
Etretinate
Gold salts
Hexachlorophene
Lovastatin
St John's Wort
Gmethylcoumarin (used in
perfumes, lotions, etc)

Note: A period of 2 weeks must pass after your last dose before your laser treatment can be performed. **Please first consult with your doctor before stopping any medication.**